

19. We adopt the positive direction choices used in the textbook so that equations such as Eq. 4-22 are directly applicable. The initial velocity is horizontal so that $v_{0y} = 0$ and $v_{0x} = v_0 = 161$ km/h. Converting to SI units, this is $v_0 = 44.7$ m/s.
- (a) With the origin at the initial point (where the ball leaves the pitcher's hand), the y coordinate of the ball is given by $y = -\frac{1}{2}gt^2$, and the x coordinate is given by $x = v_0t$. From the latter equation, we have a simple proportionality between horizontal distance and time, which means the time to travel half the total distance is half the total time. Specifically, if $x = 18.3/2$ m, then $t = (18.3/2)/44.7 = 0.205$ s.
 - (b) And the time to travel the next $18.3/2$ m must also be 0.205 s. It can be useful to write the horizontal equation as $\Delta x = v_0\Delta t$ in order that this result can be seen more clearly.
 - (c) From $y = -\frac{1}{2}gt^2$, we see that the ball has reached the height of $-\frac{1}{2}(9.8)(0.205)^2 = -0.205$ m at the moment the ball is halfway to the batter.
 - (d) The ball's height when it reaches the batter is $-\frac{1}{2}(9.8)(0.409)^2 = -0.820$ m, which, when subtracted from the previous result, implies it has fallen another 0.615 m. Since the value of y is not simply proportional to t , we do not expect equal time-intervals to correspond to equal height-changes; in a physical sense, this is due to the fact that the initial y -velocity for the first half of the motion is not the same as the "initial" y -velocity for the second half of the motion.